


BITEMARK: Traceable Multimodal Reasoning for Structured Orthodontic Reporting

Yoonha Eo

Department of Artificial Intelligence,
College of Computing, Yonsei University




- I. Background and Motivation**
- II. BITEMARK Framework Overview**
- III. Intraoral Photograph Analysis**
- IV. Intraoral Scan Analysis**
- V. Task-Level Evaluation and Method Selection**
- VI. Structured Findings and Report Assembly**
- VII. Q&A**



<https://aidentglobal.com/event>
<https://dexis.com/en-ae/orthopantomograph-op-3d-ex>
<https://www.rsipvision.com/dental-segmentation/>

Barriers to Clinical Diagnosis



**Despite the progress of deep learning in dentistry,
major challenges remain before these systems can be reliably used for clinical diagnosis.**


What's holding AI back?

Lack of Explainability

- **Black-box Decision Making**
- **Opaque Reasoning**
- **Hidden Representations**
- **Weak Feature Attribution**
- **Uncertain Causality**
- **Poor Traceability**
- **Limited Interpretability**
- **Faithfulness Gap**
- **Grounding Gap**
- **Post-hoc Explanation**
- **Spurious Correlation**
- **Unverifiable Evidence**
- **Low Clinical Transparency**
- **.....**


Lack of Explainability

- Black-box Decision Making
- Opaque Reasoning
- Hidden Representations
- Weak Feature Attribution
- Uncertain Causality
- Poor Traceability
- Limited Interpretability
- Faithfulness Gap
- Grounding Gap
- Post-hoc Explanation
- Spurious Correlation
- Unverifiable Evidence
- Low Clinical Transparency
-



Unknown Decision Logic

Input



Vision-Language Model

Output


REPORT



The patient presents on the right a full Class II molar and canine relationship, while on the left a full Class II canine relationship is present (molar relationship not definable due to the absence of tooth 26). The transverse relationships appear within normal limits; no crossbites are present. Overjet is markedly increased, overbite is within normal limits. The midlines are deviated. In the lower arch, wide diastemas are present.

Unknown Decision Logic

Input



Single Multimodal Model

Output



REPORT



The patient presents on the right a full Class II molar and canine relationship, while on the left a full Class II canine relationship is present (molar relationship not definable due to the absence of tooth 26). The transverse relationships appear within normal limits; no crossbites are present. Overjet is markedly increased, overbite is within normal limits. The midlines are deviated. In the lower arch, wide diastemas are present.

Unknown Decision Logic

Input



Output


REPORT



The patient presents on the right a full Class II molar and canine relationship, while on the left a full Class II canine relationship is present (molar relationship not definable due to the absence of tooth 26). The transverse relationships appear within normal limits; no crossbites are present. Overjet is markedly increased, overbite is within normal limits. The midlines are deviated. In the lower arch, wide diastemas are present.

Unknown Decision Logic

Input



Multimodal Task

Output


REPORT




The patient presents on the right a full Class II molar and canine relationship, while on the left a full Class II canine relationship is present (molar relationship not definable due to the absence of tooth 26). The transverse relationships appear within normal limits; no crossbites are present. Overjet is markedly increased, overbite is within normal limits. The midlines are deviated. In the lower arch, wide diastemas are present.

Traceable Multimodal Pipeline


BITEMARK: Traceable Multimodal Pipeline




Traceable Multimodal Pipeline: Intraoral Photographs



Midline: Intraoral Photographs




Midline



**Teeth (#11,21,31,41) mask centers define the two midlines.
Displacement is normalized by mean incisor width.**

Range	Midline Classification
$D < 0.063$	Centered
$0.063 \leq D < 0.16$	Slightly Deviated
$D \geq 0.16$	Deviated

Crowding: Intraoral Photographs





Crowding: Intraoral Photographs





Upper




Lower




— Smooth reference arch
— Centroid polyline
○ Top-3 excess turns




Traceable Multimodal Pipeline



Traceable Multimodal Pipeline : Intraoral Scans



Traceable Multimodal Pipeline : Intraoral Scans



Molar Relationship : Intraoral Scans

Molar Relationship




Class I




Upper and lower posterior surfaces show the selected Class I alignment.

Class II edge-to-edge



An intermediate sagittal alignment.

Class II full



Upper molar is shifted toward the anterior lower teeth.


Class III



Upper molar is shifted toward the posterior lower teeth.

1. Start with the STL tooth surface


Each triangular face supplies three surface samples



Upper first molar

2. Construct the three interior points


Place each point halfway from a vertex to the centroid



Green point: triangle centroid

3. Select the upper-molar query band


Retain points at or below the area-weighted Q65 height



Blue: selected query points Gray: excluded points


4. Find the nearest opposing surface

For each upper query point, search lower 45, 46, and 47



Upper 16: query tooth Lower 45, 46, 47: opposing surfaces

Traceable Multimodal Pipeline : Intraoral Scans




Canine Relationship : Intraoral Scans

Canine Relationship




Class II




Upper canine tip lies anterior to the lower canine-premolar reference.

Class I




Upper canine tip is near the lower canine-premolar reference.

Class III



Upper canine tip lies posterior to the lower canine-premolar reference.

Traceable Multimodal Pipeline : Intraoral Scans




Anterior Bite : Intraoral Scans

Anterior Bite




Open bite




A vertical gap separates the upper and lower incisal edge bands.

Inverted bite




Lower incisors project ahead of the upper incisors.

Normal bite




Incisal overlap is present without excessive vertical coverage.

Deep bite



The upper incisors cover a larger vertical portion of the lower incisors.

Traceable Multimodal Pipeline : Intraoral Scans



Transverse Bite : Intraoral Scans

Transverse Bite



Cross bite



Lingual



Buccal

The upper posterior tooth sits farther inward than the lower tooth.

Normal




Lingual



Buccal

The upper tooth lies slightly outward while the two transverse intervals overlap.

Scissor bite




Lingual



Buccal


The upper tooth sits far outward with little transverse interval overlap.

Traceable Multimodal Pipeline : Intraoral Scans




Curve of Spee/Wilson & Overjet : Intraoral Scans -> PointTransformer V3

FDI-Labeled Point Cloud



Registered Dual Arch Point Cloud



Task-level Results for Method Selection

Finding	PTv3 A/F1	Alternative A/F1	Selected route
Right sagittal	.750/.715	.833/.823	IOS geometry
Left sagittal	.700/.628	.833/.823	IOS geometry
Anterior bite	.600/.460	.915/.936	IOS geometry
Transverse bite	.750/.476	.880/.817	IOS geometry
Midline	.450/.437	.809/.801	IOP geometry
Curve of Spee	.719/.698	.657/.587	PTv3
Curve of Wilson	.677/.676	.573/.574	PTv3

Structured Findings : Over 28 Classes

Template

Overbite: Normal | Increased | Decreased | Open bite

Crowding (upper/lower):

None-to-mild | Mild/moderate | Moderate/severe

Molar relationship (right/left):

Class II full | Class II edge-to-edge | Class I | Class III

Canine relationship (right/left):

Class I | Class II | Class III | Not assessable

Overjet: Normal | Deep | Negative

Dental midline: Centered | Slightly deviated | Deviated

Transverse bite: Normal | Cross bite | Scissor bite

Lower curve of Spee: Normal | Increased

Lower curve of Wilson: Normal | Increased

Missing teeth:

Absent FDI identifiers among 28 permanent teeth, excluding third molars

Deterministic Report Assembly

Structured findings

```
summary_rows = [  
  {"item": "Molar relationship" ,  
   "view": "right" ,  
   "result": "Class II Full" },  
  
  {"item": "Canine relationship" ,  
   "view": "right" ,  
   "result": "Class II" },  
  
  {"item": "Midline" ,  
   "view": "front" ,  
   "result": "Deviated" },  
  
  ...  
]
```



Rule-based assembly

Right molar : Full Class II

Right canine : Class II



A [**full Class II**] molar relationship
and a [**Class II**] canine relationship
on the right.

Midline: Deviated



The dental midlines are [**deviated**]
relative to each other.



Deterministic Report Assembly

Structured findings

```
summary_rows = [  
  {"item": "Molar relationship" ,  
   "view": "right" ,  
   "result": "Class II Full" },  
  
  {"item": "Canine relationship" ,  
   "view": "right" ,  
   "result": "Class II" },  
  
  {"item": "Midline" ,  
   "view": "front" ,  
   "result": "Deviated" },  
  
  ...  
]
```

Rule-based assembly

Right molar : Full Class II

Right canine : Class II



A [full Class II] molar relationship
and a [Class II] canine relationship
on the right.

Midline: Deviated



The dental midlines are [deviated]
relative to each other.




REPORT



The patient presents on the right a full Class II molar and canine relationship, while on the left a full Class II canine relationship is present (molar relationship not definable due to the absence of tooth 26). The transverse relationships appear within normal limits; no crossbites are present. Overjet is markedly increased, overbite is within normal limits. The midlines are deviated. In the lower arch, wide diastemas are present.

BITEMARK: Traceable Multimodal Pipeline



Q & A

Contact Info

Yoonha Eo

yoonhaeo@yonsei.ac.kr / +82-10-4555-3309